2nd INTERNATIONAL CONFERENCE ON NATURAL PRODUCTS UTILIZATION:

FROM PLANTS TO PHARMACY SHELF

Plovdiv 2015



14–17 October 2015

Plovdiv, BULGARIA




2nd INTERNATIONAL CONFERENCE ON
NATURAL PRODUCTS UTILIZATION:
From Plants to Pharmacy Shelf

14–17 October 2015

BOOK OF ABSTRACTS

Plovdiv • Bulgaria

The 2nd International Conference on Natural Products Utilization: from Plants to Pharmacy Shelf (14–17 October, 2015), Plovdiv (Bulgaria) is organized with the financial support of the Ministry of Science and Education, Republic of Bulgaria



МИНИСТЕРСТВО НА ОБРАЗОВАНИЕТО, МЛАДЕЖТА И НАУКАТА

Joint meeting with the Phytochemical Society of Europe and Bulgarian Phytochemical Society





Illustration: Cover photo (*Haberlea rhodopensis*) provided courtesy of I. Aneva
Editor: Milen I. Georgiev, PhD

2nd INTERNATIONAL CONFERENCE ON NATURAL PRODUCTS UTILIZATION: From Plants to Pharmacy Shelf

14–17 October 2015, Plovdiv, Bulgaria

CHAIRS

Milen I. GEORGIEV
Institute of Microbiology,
Bulgaria

Kalina I. ALIPIEVA
Institute of Organic Chemistry
with Centre of Phytochemistry
Bulgaria

HONORARY CHAIRPERSON

Vassya BANKOVA
Corresponding member of BAS
Institute of Organic Chemistry with Centre of Phytochemistry, Bulgaria

SCIENTIFIC COMMITTEE MEMBERS

Bharat B. AGGARWAL
The University of Texas M. D. Anderson
Cancer Center (USA)
Strahil BERKOV
Institute of Biodiversity and
Ecosystem Research
(Bulgaria)
Carlos L. CESPEDES
University of Bio Bio (Chile)
Tossaton CHAROONRATANA
Rangsit University (Thailand)
Marc DIEDERICH
Seoul National University (Korea)
Vladimir DIMITROV
IOCCP (Bulgaria)
Petya DIMITROVA
Institute of Microbiology (Bulgaria)
Albena DINKOVA-KOSTOVA
University of Dundee (UK)/
The Johns Hopkins University (USA)
Balik DZHAMBAZOV
University of Plovdiv (Bulgaria)
Tsanko GECHEV
University of Potsdam (Germany)
Vasil GEORGIEV
Florida A & M University (USA)
Elvira GILLE
National Institute of R&D for
Biological Sciences (Romania)
Ilkay Erdogan ORHAN
Gazi University (Turkey)
Ilza PAJEVA
Corresponding member of BAS
Institute of Biophysics and
Biomedical Engineering (Bulgaria)
Javier PALAZON
University of Barcelona (Spain)
Milena POPOVA
IOCCP (Bulgaria)
Victoria SARAFIAN
Medical University–Plovdiv
(Bulgaria)
Pavleta SHESTAKOVA
IOCCP (Bulgaria)

Stoyan A. SHISHKOV
Sofia University (Bulgaria)
Svetlana SIMOVA
Institute of Organic Chemistry with Centre of Phytochemistry (Bulgaria)
Krystyna SKALICKA-WOZNIAK
Medical University of Lublin (Poland)
Leandros SKALTSOUNIS
University of Athens (Greece)
Gjoshe STEFKOV
University Ss. Cyril and Methodius (Macedonia)
Zora Dajić STEVANOVIC
University of Belgrade (Serbia)
Iliana IONKOVA
Medical University of Sofia (Bulgaria)
Diana IVANOVA
Medical University of Varna (Bulgaria)
Veneta KAPCHINA-TOTEVA
Sofia University (Bulgaria)
Spiro M. KONSTANTINOV
Medical University of Sofia (Bulgaria)
Ilina KRASTEVA
Medical University of Sofia (Bulgaria)

Jutta LUDWIG-MÜLLER
TU Dresden (Germany)
Adam MATKOWSKI
Medical University of Wroclaw (Poland)
Albena STOYANOVA
University of Food Technologies (Bulgaria)
Miroslav STRNAD
Chairman of the PSE, Institute of Experimental Botany (Czech Republic)
Robert VERPOORTE
Leiden University (The Netherlands)
Alvaro M. VILJOEN
Tshwane University of Technology (South Africa)
Jean-Luc WOLFENDER
University of Geneva (Switzerland)
Jianbo XIAO
Macau University (PR China)
Ning-Sun YANG
Academia Sinica (ROC)
Danijela MISIC
Institute for Biological Research Siniša Stanković (Serbia)

ORGANIZING COMMITTEE MEMBERS

Andrey MARCHEV – Chair
Zhenya YORDANOVA
Snezhana RUSINOVA-VIDEVA
Elka GENOVA
Katerina GEORGIEVA
Georgi ZAHMANOV
Tsvetanka TENEVA-ANGELOVA
Lidiya GEORGIEVA

PP 153

IMMUNOLOGICALLY ACTIVE POLYSACCHARIDE-CONTAINING COMPLEXES FROM TRADITIONAL AND PERSPECTIVE EUROPEAN MEDICINAL PLANTS

Yordan Georgiev[1], Manol Ognyanov[1], Irina Yanakieva[1], Veselin Kussovski[3], Hiroaki Kiyohara[4], Petko Denev[1], Daniela Antonova[2], Milan Čiž[5], Antonín Lojek[5], Haruki Yamada[4], Maria Kratchanova[1]

[1] *Laboratory of Biologically Active Substances, Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, Plovdiv, Bulgaria*
[2] *Laboratory of Experimental Chromatography and Mass Spectrometry, Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, Sofia, Bulgaria*
[3] *Department of Infectious Microbiology, Stephan Angeloff Institute of Microbiology, Bulgarian Academy of Sciences, Sofia, Bulgaria*
[4] *Department of Drug Discovery Sciences, Kitasato Institute for Life Sciences, Kitasato University, Tokyo, Japan*
[5] *Department of Free Radical Pathophysiology, Institute of Biophysics, Czech Academy of Sciences, Brno, Czech Republic*

In the present study we investigated the chemical composition and immunomodulating activity of crude water-soluble polysaccharide mixtures extracted with boiling water from linden (flower), thyme (aerial part), nettle (leaf and root), lavender (aerial part), mountain tea (aerial part), common purslane (aerial part) and soapwort (aerial part). Water-soluble polysaccharides presented in relatively low amounts (2.3-8.6 %) in the studied herbs. All herbs contained mixtures of acidic and neutral (hetero)polysaccharides with the following monosaccharides: Ara, Gal, Rha, Xyl, Man, Glc, Fuc, GalA and GlcA. The total sugars varied between 46.1-77.6 %, as anhydrouronic acid content was the predominant part for most polysaccharide mixtures, except for these from nettle (root) and soapwort. The molecular weight of polysaccharides was between 7.6 - 1950 kDa. The presence of proteins, polyphenols and saponins was also estimated. All polysaccharide mixtures, except that from mountain tea showed high *in vitro* anti-complementary activity (82.6-49.8 %) at 78 μg/mL expressed as total complement hemolysis of target rabbit erythrocytes. Some samples exhibited *in vitro* moderate to high activity against murine intestinal Peyer's patch immunocompetent cells. Activation of *in vitro* ROS production from human whole blood neutrophils was observed for samples with reduced content of lower molecular weight constituents. Additionally, higher polyphenol content in some crude polysaccharides was related to moderate *in vitro* antioxidant activity. The most potent immunomodulating polysaccharide mixtures were isolated from nettle (root) and common purslane.

Acknowledgements: This work was supported by project BG161PO0003-1.1.05-0024-C0001, OP "Development of the Competitiveness of Bulgarian Economy" and project BG051PO001/3.3-05-0001, OP "Human Resource Development".

PP 57

VASOPROTECTIVE PROPERTIES OF ARONIA MELANOCARPA – A HISTOLOGICAL AND MORPHOMETRIC STUDY

Elena Daskalova[1], Slavi Delchev[1], Ilia Bivolarski[2], Petko Denev[3,4], Maria Kratchanova[3,4], Petar Cvetkov[5], Melek Kaluch[5]

[1] Deparment of Anatomy, Histology and Embryology, Medical University of Plovdiv, Bulgaria
[2] Deparment of general and clinical pathology, Medical University of Plovdiv, Bulgaria
[3] Laboratory of Biologically Active Substances, Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, Plovdiv, Bulgaria.
[4] ITC-Innovative-Technological Centre Ltd., Plovdiv, Bulgaria
[5] Student of medicine, Medical University of Plovdiv, Plovdiv, Bulgaria

The social significance of age-related diseases is determined by their global role in mortality and morbidity, particularly in economically developed countries. The most severe forms of coronary heart disease and cerebrovascular disease are the main reasons for lethality, whereas the "milder" forms often lead to disability of patients in working age. Changes in elastic and muscular arteries walls, resulting from age-related restructuring and progression of atherosclerotic lesions, underlie these diseases. Their prevention through administration of natural products is a research area with huge potential, and application of natural antioxidants is one of the leading strategies to retard vascular aging. *Aronia melanocarpa* juice is a rich source of polyphenols and is characterized by very high antioxidant activity *in vitro*. The aim of the current study was to investigate the effect of aronia juice intake on age-related vascular changes of rat aortic walls. We used a model of aging male rats, whose thoracic aorta walls were subjected to macroscopic, histopathological (hematoxilyn-eosin), histochemical (azan and orcein) and morphometric studies. The comparative analysis between the target group of old animals supplemented with aronia juice; young untreated rats and old controls (not supplemented), revealed that aronia-supplemented animals were characterized with reduced atherosclerotic lesions and a lower level of restructuring of aortic walls. These data confirm that *Aronia melanocarpa* juice successfully retards age-related vascular aging, and can be recommended as a prophylactic tool for healthy aging.

Acknowledgements: This work was funded by grant BG161PO003-1.1.05-0024 of Operational Programme "Development of the Competitiveness of Bulgarian Economy".