



Honorary Chair

Prof. Dr. Erhan TABAKOĞLU (Trakya University Rector)

Chair

Prof.Dr. Mesut KAÇAN (Head of Chemistry Department-Trakya University)

Co-Chair

Assoc.Prof.Dr. Hakkı M. ÖZCAN


Assoc.Prof.Dr. Mesut BOZ

Asst.Prof.Dr. Kenan SEZER


Congress Secretary

Assoc.Prof.Dr. H.R.Ferhat KARABULUT

Assoc.Prof.Dr. Murat TURKYILMAZ



Scientific Committee



Prof.Dr. Skender Topi	(UNIEL)
Prof.Dr. Nikolai D. Denkov	(SOFIA UNI.)
Prof.Dr. Nikos Hadjichristidis	(KAUST)
Prof.Dr. Metin Balcı	(METU)
Prof.Dr. Cihangir Tanyeli	(METU)
Prof.Dr. Yusuf Yağcı	(ITU)
Prof.Dr. Vassil Borrisov Delchev	(PLOVDIV UNI.)
Prof.Dr. Şefik Süzer	(BILKENT UNI.)
Prof.Dr. Agim Shabani	(TETOVA UNI.)
Prof.Dr. Mustafa M. Demir	(IIT)
Prof.Dr. Athanasios G. Mamalis	(DEMOCRITUS UNI.)
Prof.Dr. Ahmet Çolak	(KTU)
Prof.Dr. Ömer Zaim	(TRAKYA UNI.)
Prof.Dr. Mustafa Sözbilir	(ATATÜRK UNI.)
Prof.Dr. Kleomenis Barlos	(CBL)
Prof.Dr. Amira Cipukovic	(TUZLA UNI.)
Prof.Dr. Trajce Stafilov	(UKIM)
Prof.Dr. Panagiotis Spathis	(AUT)
Prof.Dr. Sotir Mali	(UNIEL)
Assoc.Prof.Dr. Jehad Almaliti	(JORDAN UNI.)
Assoc.Prof.Dr. Kiril Gavazov	(MUP)

OrganizationCommittee

Prof.Dr. Hülya YAĞAR

Prof.Dr. Mesut KACAN

Prof.Dr. Nesrin BEYNEK

Prof.Dr. Ömer ZALM

Prof.Dr. Yasemin BAKIRCIOĞLU KURTULUŞ

Prof.Dr.Yüksel BAYRAK

Assoc.Prof.Dr. Çiğdem BATIGÖÇ

Assoc.Prof.Dr. Dilek BAKIRCIOĞLU

Assoc.Prof.Dr. Eylem BAYIR

Assoc.Prof.Dr. H.R.Ferhat KARABULUT

Assoc.Prof.Dr. Hakkı M. ÖZCAN

Assoc.Prof.Dr. Mesut BOZ

Assoc.Prof.Dr. Murat TURKYILMAZ

Assoc.Prof.Dr. Özlen ALTUN

Assoc.Prof.Dr. Şebnem Selen İŞBİLİR

Asst.Prof.Dr. Gühergöl ULUÇAM

Asst.Prof.Dr. Hasan ÖZYILDIRIM

Asst.Prof.Dr. Hayrettin BEYNEK

Asst.Prof.Dr. Hüsnüye DURMAZ

Asst.Prof.Dr. Kenan SEZER

Asst.Prof. Dr. Taliha SİDİM

Asst.Prof.Dr. Yıldız KALEBAŞI

Dr. Ayşen ŞUEKİNCİ YILMAZ

Dr. Birgül KARADAĞ

Dr. Hafize ÖZCAN

Dr. Zuhal HOŞGÖR

Dr. Ali Osman KARATAVUK



(M25-PS4-20185) Desing, synthesis and antitrichinellosis activity of 1,3-disubstituted benzimidazol-2-ones

Oisaemi Izevbekhai¹, Kameliya Anichina-Zarkova¹, Anelia Mavrova¹, Dimitar Vutchev², Galya Popova-Daskalova², Denitsa Yancheva³, Simeon Stoyanov³

¹University Of Chemical Technology And Metallurgy

²Medical University

³Bulgarian Academy Of Sciences

kameliya_anichina@uctm.edu

Benzimidazole anthelmintic drugs as thiabendazole, mebendazole, albendazole, etc. are widely used in veterinary and human medicine. Because of their high efficacy, prolonged use of these compounds has led to the development of resistance by many helminth strains.

For this reason, the need for the design and synthesis of new benzimidazole molecules to overcome existing resistance has increased.

Novel 1,3-disubstituted benzimidazol-2-ones were prepared using a multi-step synthetic approach that started from 4-substituted-1,2-diaminobenzenes. The structures of the targeted compounds were confirmed by IR and ¹H NMR spectra. In addition, the molecular geometry and electron structure of these molecules were theoretically evaluated using density functional theory (DFT) methods.

The compounds exhibited remarkable activity *in vitro* against isolated *Trichinella spiralis* muscle larvae.

Keywords: 1,3-disubstituted benzimidazol-2-ones, antitrichinellosis activity; DFT

References

- [1] William. C. Campbell (Ed.), *Trichinella and Trichinosis*, Plenum Press, USA and London, UK.- 1983.-P. 340-355
- [2] E. Pozio, D. Sacchini, L. Sacchi, A. Tamburrini, F. Alberici. Failure of Mebendazole in the Treatment of Humans with *Trichinella spiralis* Infection at the Stage of Encapsulating Larvae.// *Clin.Infect. Dis.*- 2001. V. 32.-P. 638-642
- [3] Anelia Ts. Mavrova, Kamelya K. Anichina, Dimitar I. Vutchev, Jordan A. Tsenov, Magdalena S. Kondeva, Mitka K. Micheva. Synthesis and antitrichinellosis activity of some 2-substituted-[1,3]thiazolo[3,2-a]benzimidazol-3(2H)-ones.// *Bioorg. and Med. Chem.* -2005. V. 13. -P. 5550-5559
- [4] Anelia Ts. Mavrova, Kamelya K. Anichina, Dimitar I. Vutchev, Jordan A. Tsenov, Pavletta S. Denkova, Magdalena S. Kondeva, Mitka K. Micheva. Antihelminthic activity of some newly synthesized 5(6)-(un)substituted-1H-benzimidazol-2-ylthioacetyl piperazine derivatives. *Eur. J. Med.Chem.*// 2006. V. 41(12).-P. 1412-1420



(M26-PS4-20190) DFT and experimental IR study on the conversion of 6(7)-nitro-[1,3]thiazolo[3,2-a]-benzimidazole-3(2H)-ones with antiparasitic activity into anion products

Denitsa Yancheva¹, Kameliya Anichina-Zarkova², Simeon Stoyanov¹, Anelia Mavrova², Dimitar Vutchev³, Galya Popova-Daskalova³

¹ Bulgarian Academy Of Sciences

² University Of Chemical Technology And Metallurgy

³ Medical University

kameliya_anichina@uctm.edu

Nitroheterocyclic drugs ability to generate free radicals under anaerobic growth conditions, thus affecting multiple cellular pathways, has been is a successful strategy to develop effective drugs for treatment of parasitic worm infestations. Nitroheterocyclic medications are widely used to cure giardiasis, trichuriasis, filariasis, neurocysticercosis, hydatid disease, pinworm disease, ascariasis, Chagas disease and African sleeping sickness [1,2].

As part of our research on antiparasitic benzimidazole derivatives, various groups of benzimidazole derivatives were synthesized and evaluated for their activity *in vitro* and *in vivo* against different protozoa. The 2-arilyden-thiazolo[3,2-a]benzimidazol-3(2H)-ones containing different substituents at the 6(7)-position showed remarkable activity against the intestinal and muscle phases of *Trichinella spiralis* in white mice [3]. Having in mind the key role of the nitro group for manifestation of antiparasitic activity, recently we focused our attention on characterizing the metabolites resulting from bioreduction of nitrobenzimidazoles into more details. The possible conversion of nitro compounds into radical anion species can be conveniently studied by electrochemical reduction coupled with IR spectroscopy measurements and quantum chemical calculations on the structure, vibrational spectra and ability of the molecules to accept electrons [4]. Herein we report the results on the electrochemical reduction of some representative 6(7)-nitro-[1,3]thiazolo[3,2-a]-benzimidazole-3(2H)-one derivatives. Molecular structure, electronic charge distribution, vibrational spectra and molecular parameters related to nitro reduction were studied by DFT calculations.

Keywords: Nitrobenzimidazoles, antiparasitic activity, IR study, DFT

References

- [1] S. Patterson, S. Wyllie, Nitro drugs for the treatment of trypanosomatid diseases: past, present, and future prospects, Trends Parasitol. 2014, 30, 289- 298.
- [2] A.M. Jarrad, A. Debnath, Y. Miyamoto, K.A. Hansford, R. Pelingon, M.S. Butler, T. Bains, T. Karoli, M.A.T. Blaskovich, L. Eckmann, M.A. Coopera, Nitroimidazole carboxamides as antiparasitic agents targeting Giardia lamblia, Entamoeba histolytica and Trichomonas vaginalis, Eur. J. Med. Chem. 2016, 120, 353–362.
- [3] A. Ts. Mavrova, K. K. Anichina, D. I. Vutchev, J. A. Tsenov, M. S. Kondeva, M. K. Micheva, Bioorg. Med. Chem. 2005, 13, 5550- 5559.
- [4] D. Yancheva, S. Stoyanov, N. Anastassova, A. Ts. Mavrova, IR study on the electrochemical generation of a nitro radical anion by a hepatotoxic N,N'-disubstituted benzimidazole-2-thione, Vibr. Spectrosc. 2017, 92, 200–214.