

# 24<sup>th</sup> Congress of Chemists and Technologists of Macedonia

## BOOK of ABSTRACTS



11-14 September 2016  
Ohrid, Republic of Macedonia

MPCE 021

**MOLECULAR DOCKING STUDY ON BINDING MODES OF  
ISOXAZOLO- AND THIAZOLOHYDRAZILIDENE-  
CHROMANE-2,4-ONES WITH B-RAF KINASE**

Denitsa Yancheva<sup>1</sup>, Bozhanka Mikhova<sup>1</sup>, Lulzime Ballazhi<sup>2</sup>, Miroslav Rangelov<sup>1</sup>, Faik Imeri<sup>3</sup>, Emil Popovski<sup>4</sup>, Ahmed Jashari<sup>5</sup>, Goran Stojkovikj<sup>4</sup>, Kristina Mladenovska<sup>2</sup>

e-mail: krml@ff.ukim.edu.mk

- 1- Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, Acad. G. Bonchev Str., build. 9, 1113 Sofia, Bulgaria
- 2- Faculty of Pharmacy, Center of Biomolecular Pharmaceutical Analyses, University "Ss Cyril and Methodius", Mother Theresa 47, 1000 Skopje, Republic of Macedonia
- 3- Institute of Physiology, University of Zürich, CH-8057, Zürich, Switzerland
- 4- Institute of Chemistry, Faculty of Natural Sciences and Mathematics, University "Ss. Cyril and Methodius", Arhimedova Str. No. 3, PO Box 162, 1000 Skopje, Republic of Macedonia
- 5- Faculty of Natural Sciences & Mathematics, State University of Tetovo, Pashe Deralla bb, 1200 Tetovo, Republic of Macedonia

Recently a series of novel isoxazolo- and thiazolohydrazilidene-chromane-2,4-ones was synthesized and shown to exhibit cytotoxicity on human breast cancer cell line MDA-MB-231 [1]. MDA-MB-231 cells are very sensitive to treatment with B-Raf kinase inhibitors blocking the RAF/MEK/ERK signaling pathway and affecting the tumor growth. Among the three RAF isoforms in humans (A-Raf, B-Raf, and C-Raf), B-Raf is the most critical to mediate Ras activity. A significant fraction of melanoma, colorectal, thyroid and breast cancers have activating B-Raf mutations, particularly at valine 599 [2].

The results of a molecular docking study conducted in effort to throw more light on the possible interaction of the synthesized isoxazolo- and thiazolohydrazilidene-chromane-2,4-ones with B-Raf are presented. The compounds were docked into the crystal structure of human oncogenic <sup>V599E</sup>B-Raf in complex with the inhibitor sorafenib. All of them occupied an internal pocket formed by the activation segment and the P-loop of <sup>V599E</sup>B-Raf. The ligand-pocket binding was driven by hydrogen bonding to amino acid residues Lys482, Glu500, Thr528 and Asp593 and hydrophobic interactions with Leu504, Leu513, Gly592 and Phe594. In this way, the ligands connected simultaneously to the activation segment and the catalytic loop of <sup>V599E</sup>B-Raf, were being able to promote an inactive conformation of the enzyme and hinder its phosphorylation.

**Key words:** hydrazinyldiene-chroman-2,4-diones, molecular docking, B-Raf kinase

**References:**

- [1] Jashari A, Imeri F, Ballazhi L, Shabani A, Mikhova B, Dräger G, Popovski E, Huwiler A (2014) Synthesis and cellular characterization of novel isoxazolo- and thiazolohydrazinyldiene-chroman-2,4-diones on cancer and non-cancer cell growth and death. *Bioorg Med Chem* 22:2655-2661.
- [2] M. J. Garnett and R. Marais, Guilty as charged: B-RAF is a human oncogene, *Cancer Cell* 2004, 6, 313.