



KARDIONEfrologija CARDIONEPHROLOGY





KARDIONEfrologija

CARDIONEPHROLOGY

3rd International Cardioneurology and Hypertension Congress
"Cardioneurology Up To Date"

3rd International Cardioneurology and Hypertension Congress
"Cardioneurology Up To Date"

Organisers:

Prof. Dr. Sonja Radenković
Prof. Dr. Andrija Šmelcerović

Reviewers:

Prof. Dr. Werner Siems
Prof. Dr. Gerard London
Prof. Dr. Zorana Vasiljević

Computer processing and design:

Katarina Tomović
Marija Predić

Translator:

Marija Slavković

Editors:

Prof. Dr. Sonja Radenković
Prof. Dr. Andrija Šmelcerović

Printing / Publishing:

Punta – Niš

Niš, 2017

ISBN 978-86-7990-174-3

2-Imino-benzimidazoles and 1,3-thiazolo[3,2-a]benzimidazolones as xanthine oxidase inhibitors

Andrija Šmelcerović¹, Žaklina Šmelcerović², Katarina Tomović³, Gordana Kocić⁴, Denitsa Yancheva⁵, Anelia Ts. Mavrova⁶

¹Department of Chemistry, Faculty of Medicine, University of Nis, Bulevar Dr Zorana Djindjica 81, 18000 Nis, Serbia

²Center for Biomedicinal Science, Faculty of Medicine, University of Nis, Bulevar Dr Zorana Djindjica 81, 18000 Nis, Serbia

³Department of Pharmacy, Faculty of Medicine, University of Nis, Bulevar Dr Zorana Djindjica 81, 18000 Nis, Serbia

⁴Institute of Biochemistry, Faculty of Medicine, University of Nis, Bulevar Dr Zorana Djindjica 81, 18000 Nis, Serbia

⁵Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Science, Acad. G Bonchev Str., build. 9, 1113 Sofia, Bulgaria

⁶University of Chemical Technology and Metallurgy, 8 Kliment Ohridski Blvd., 1756 Sofia, Bulgaria

Xanthine oxidase is flavoprotein enzyme that catalyzes oxidation of hypoxanthine to xanthine, and then xanthine to uric acid. The concentration of xanthine oxidase in serum is increased in pathological conditions such as hepatitis, inflammation, ischemia-reperfusion injury, carcinogenesis and aging. Allopurinol is a known competitive inhibitor of xanthine oxidase widely used in the therapy of gout. There is a need for new xanthine oxidase inhibitors, which would be approximately as efficient as allopurinol, and thereby show fewer side effects. Benzimidazole heterocycle system is an important pharmacophore in medicinal chemistry. Its derivatives show different biological activity.

In this study, five 2-imino-benzimidazoles and four 1,3-thiazolo[3,2-a]benzimidazolones were evaluated for inhibitory activity against commercial enzyme xanthine oxidase. Additionally, an *in silico* study of pharmacokinetic and toxicological properties of the most potent inhibitors was performed.

Commercial bovine milk xanthine oxidase was used for *in vitro* test of enzyme inhibition which was based on the spectrophotometric determination of uric acid formation at 293 nm. Allopurinol was used as a positive control. Pharmacokinetic and toxicological properties of studied benzimidazoles were calculated using OSIRIS Property Explorer, admetSAR and Toxtree.

Four of nine examined benzimidazoles showed an inhibitory effect on xanthine oxidase with an IC₅₀ value below 150 μM. The highest inhibitory effect on xanthine oxidase showed compound 5-methyl-1,3-bis(3-phenylpropyl)-1*H*-benzo[d]imidazol-2(3*H*)-imine hydrobromide.

2-Imino-benzimidazoles and 1,3-thiazolo[3,2-a]benzimidazolones are novel non-purine inhibitors of xanthine oxidase.

**Inhibicija ksantin-oksidade 2-imino-benzimidazolima i
1,3-tiazolo[3,2-a]benzimidazolona**

Andrija Šmelcerović¹, Žaklina Šmelcerović², Katarina Tomović³, Gordana Kocić⁴, Denitsa Yancheva⁵, Anelia Ts. Mavrova⁶

¹Katedra Hemija, Medicinski fakultet, Univerzitet u Nišu, Bulevar dr Zorana Đinđića 81, 18000 Niš, Srbija

²Naučnoistraživački centar za biomedicinu, Medicinski fakultet, Univerzitet u Nišu, Bulevar dr Zorana Đinđića 81, 18000 Niš, Srbija

³Katedra Farmacija, Medicinski fakultet, Univerzitet u Nišu, Bulevar dr Zorana Đinđića 81, 18000 Niš, Srbija

⁴Katedra Biohemija, Medicinski fakultet, Univerzitet u Nišu, Bulevar dr Zorana Đinđića 81, 18000 Niš, Srbija

⁵Institut za organsku hemiju sa centrom za fitohemiju, Bugarska akademija nauka, Acad. G Bonchev Str., build. 9, 1113 Sofija, Bugarska

⁶Univerzitet za hemijsku tehnologiju i metalurgiju, 8 Kliment Ohridski Blvd., 1756 Sofija, Bugarska

Ksantin-oksida je flavoproteinski enzim koji katalizuje oksidaciju hipoksantina u ksantin, a zatim ksantina u mokraćnu kiselinu. Koncentracija ksantin-oksidade u serumu povećana je u patološkim stanjima kao što su hepatitis, zapaljenje, ishemija-reperfuzija, karcinogeneza i starenje. Alopurinol je poznati kompetitivni inhibitor ksantin-oksidade široko zastupljen u terapiji gihta. Postoji potreba za novim inhibitorima ksantin-oksidade koji bi bili približno efikasni kao alopurinol, a pritom pokazivali manje sporednih efekata od njega. Benzimidazolni heterociklusni sistem je važna farmakofora u farmaceutskoj hemiji. Derivati benzimidazola pokazuju različitu biološku aktivnost.

U ovom radu je ispitan uticaj pet 2-imino-benzimidazola i četiri 1,3-tiazolo[3,2-a]benzimidazolona na aktivnosti komercijalnog enzima ksantin-oksidade. Takođe, *in silico* metodama je izvršeno predviđanje farmakokinetičkih i toksikoloških parametara najefikasnijih inhibitora.

In vitro test inhibicije komercijalne ksantin-oksidade, izolovane iz govedeg mleka, se zasniva na spektrofotometrijskom određivanju dobijene mokraćne kiseline na 293 nm. Alopurinol je korišćen kao pozitivna kontrola. Farmakološki i toksikološki parametri ispitivanih benzimidazola su predviđani kompjuterskim programima OSIRIS Property Explorer, admetSAR i Toxtree.

Četiri od devet ispitivanih benzimidazola je pokazalo inhibitorni efekat na komercijalnoj ksantin-oksidazi sa IC₅₀ vrednošću manjom od 150 µM. Najjači inhibitorni efekat na ksantin-oksidazu pokazao je 5-metil-1,3-bis(3-fenilpropil)-1*H*-benzo[d]imidazol-2(3*H*)-imin hidrobromid.

2-Imino-benzimidazoli i 1,3-tiazolo[3,2-a]benzimidazoloni predstavljaju nove nepurinske inhibitore ksantin-oksidade.