


NATURAL PRODUCT COMMUNICATIONS

An International Journal for Communications and Reviews Covering all
Aspects of Natural Products Research



Volume 10. Issue 8. Pages 1333-1494. 2015
ISSN 1934-578X (printed); ISSN 1555-9475 (online)
www.naturalproduct.us

EDITOR-IN-CHIEF

DR. PAWAN K AGRAWAL

Natural Product Inc.
7963, Anderson Park Lane,
Westerville, Ohio 43081, USA
agrawal@naturalproduct.us

EDITORS

PROFESSOR ALEJANDRO F. BARRERO

Department of Organic Chemistry,
University of Granada,
Campus de Fuente Nueva, s/n, 18071, Granada, Spain
afbarre@ugr.es

PROFESSOR ALESSANDRA BRACA

Dipartimento di Chimica Bioorganica e Biofarmacia,
Università di Pisa,
via Bonanno 33, 56126 Pisa, Italy
braca@farm.unipi.it

PROFESSOR DE-AN GUO

State Key Laboratory of Natural and Biomimetic Drugs,
School of Pharmaceutical Sciences,
Peking University,
Beijing 100083, China
gda5958@163.com

PROFESSOR YOSHIHIRO MIMAKI

School of Pharmacy,
Tokyo University of Pharmacy and Life Sciences,
Horinouchi 1432-1, Hachioji, Tokyo 192-0392, Japan
mimaki@ps.toyaku.ac.jp

PROFESSOR STEPHEN G. PYNE

Department of Chemistry
University of Wollongong
Wollongong, New South Wales, 2522, Australia
spyne@uow.edu.au

PROFESSOR MANFRED G. REINECKE

Department of Chemistry,
Texas Christian University,
Forts Worth, TX 76129, USA
m.reinecke@tcu.edu

PROFESSOR WILLIAM N. SETZER

Department of Chemistry
The University of Alabama in Huntsville
Huntsville, AL 35809, USA
wsetzer@chemistry.uah.edu

PROFESSOR YASUHIRO TEZUKA

Faculty of Pharmaceutical Sciences
Hokuriku University
Ho-3 Kanagawa-machi, Kanazawa 920-1181, Japan
y-tezuka@hokuriku-u.ac.jp

PROFESSOR DAVID E. THURSTON

Department of Pharmacy and Forensic Science,
King's College London,
Britannia House, 7 Trinity Street,
London SE1 1DB, UK.
david.thurston@kcl.ac.uk

HONORARY EDITOR

PROFESSOR GERALD BLUNDEN

The School of Pharmacy & Biomedical Sciences,
University of Portsmouth,
Portsmouth, PO1 2DT U.K.
axuf64@dsl.pipex.com

ADVISORY BOARD

Prof. Viqar Uddin Ahmad
Karachi, Pakistan

Prof. Giovanni Appendino
Novara, Italy

Prof. Yoshinori Asakawa
Tokushima, Japan

Prof. Roberto G. S. Berlinck
São Carlos, Brazil

Prof. Anna R. Bilia
Florence, Italy

Prof. Maurizio Bruno
Palermo, Italy

Prof. César A. N. Catalán
Tucumán, Argentina

Prof. Josep Coll
Barcelona, Spain

Prof. Geoffrey Cordell
Chicago, IL, USA

Prof. Fatih Demirci
Eskişehir, Turkey

Prof. Ana Cristina Figueiredo
Lisbon, Portugal

Prof. Cristina Gracia-Viguera
Murcia, Spain

Dr. Christopher Gray
Saint John, NB, Canada

Prof. Dominique Guillaume
Reims, France

Prof. Duvvuru Gunasekar
Tirupati, India

Prof. Hisahiro Hagiwara
Niigata, Japan

Prof. Tsukasa Iwashina
Tsukuba, Japan

Prof. Leopold Jirovetz
Vienna, Austria

Prof. Vladimir I Kalinin
Vladivostok, Russia

Prof. Phan Van Kiem
Hanoi, Vietnam

Prof. Niel A. Koorbanally
Durban, South Africa

Prof. Chiaki Kuroda
Tokyo, Japan

Prof. Hartmut Laatsch
Gottingen, Germany

Prof. Marie Lacaille-Dubois
Dijon, France

Prof. Shoen-Sheng Lee
Taipei, Taiwan

Prof. Imre Mathe
Szeged, Hungary

Prof. M. Soledade C. Pedras
Saskatoon, Canada

Prof. Luc Pieters
Antwerp, Belgium

Prof. Peter Proksch
Düsseldorf, Germany

Prof. Phila Raharivelomanana
Tahiti, French Polynesia

Prof. Luca Rastrelli
Fisciano, Italy

Prof. Stefano Serra
Milano, Italy

Prof. Monique Simmonds
Richmond, UK

Dr. Bikram Singh
Palampur, India

Prof. John L. Sorensen
Manitoba, Canada

Prof. Johannes van Staden
Scottsville, South Africa

Prof. Valentin Stonik
Vladivostok, Russia

Prof. Winston F. Tinto
Barbados, West Indies

Prof. Sylvia Urban
Melbourne, Australia

Prof. Karen Valant-Vetschera
Vienna, Austria

INFORMATION FOR AUTHORS

Full details of how to submit a manuscript for publication in Natural Product Communications are given in Information for Authors on our Web site <http://www.naturalproduct.us>.

Authors may reproduce/republish portions of their published contribution without seeking permission from NPC, provided that any such republication is accompanied by an acknowledgment (original citation)-Reproduced by permission of Natural Product Communications. Any unauthorized reproduction, transmission or storage may result in either civil or criminal liability.

The publication of each of the articles contained herein is protected by copyright. Except as allowed under national "fair use" laws, copying is not permitted by any means or for any purpose, such as for distribution to any third party (whether by sale, loan, gift, or otherwise); as agent (express or implied) of any third party; for purposes of advertising or promotion; or to create collective or derivative works. Such permission requests, or other inquiries, should be addressed to the Natural Product Inc. (NPI). A photocopy license is available from the NPI for institutional subscribers that need to make multiple copies of single articles for internal study or research purposes.

To Subscribe: Natural Product Communications is a journal published monthly. 2015 subscription price: US\$2,595 (Print, ISSN# 1934-578X); US\$2,595 (Web edition, ISSN# 1555-9475); US\$2,995 (Print + single site online); US\$595 (Personal online). Orders should be addressed to Subscription Department, Natural Product Communications, Natural Product Inc., 7963 Anderson Park Lane, Westerville, Ohio 43081, USA. Subscriptions are renewed on an annual basis. Claims for nonreceipt of issues will be honored if made within three months of publication of the issue. All issues are dispatched by airmail throughout the world, excluding the USA and Canada.

Effects on MC3T3-E1 Cells and *In silico* Toxicological Study of Two 6-(Propan-2-yl)-4-methyl-morpholine-2,5-diones

Marija Vukelić-Nikolić^a, Ana Kolarević^b, Katarina Tomović^b, Denitsa Yancheva^c, Emiliya Cherneva^d, Stevo Najman^a, and Andrija Šmelcerović^{b, e*}

^aInstitute of Biology and Human Genetics, Faculty of Medicine, University of Nis, Bulevar Dr Zorana Djindjica 81, 18000 Nis, Serbia

^bDepartment of Pharmacy, Faculty of Medicine, University of Nis, Bulevar Dr Zorana Djindjica 81, 18000 Nis, Serbia

^cLaboratory of Structural Organic Analysis, Institute of Organic Chemistry with Centre of Phytochemistry, Bulgarian Academy of Sciences, Acad. G. Bonchev Str., build. 9, 1113 Sofia, Bulgaria

^dDepartment of Chemistry, Faculty of Pharmacy, Medical University of Sofia, 2 Dunav Str., 1000 Sofia, Bulgaria

^eDepartment of Chemistry, Faculty of Medicine, University of Nis, Bulevar Dr Zorana Djindjica 81, 18000 Nis, Serbia

a.smelcerovic@yahoo.com (or andrija.smelcerovic@medfak.ni.ac.rs)

Received: April 23rd, 2015; Accepted: May 18th, 2015

Recently, we found that two cyclodipeptides, 3,6-di-(propan-2-yl)-4-methyl-morpholine-2,5-dione (**1**) and 3-(2-methylpropyl)-6-(propan-2-yl)-4-methyl-morpholine-2,5-dione (**2**), are excellent inhibitors of xanthine oxidase. In order to obtain more information about the toxicological potential of compounds **1** and **2** on bone cells, the current study was designed to evaluate the effect of these compounds on viability and proliferation of MC3T3-E1 cells. Compound **1** showed neither cytotoxic nor stimulatory effect on cell viability, while compound **2** showed a slight stimulatory effect on cell viability. Both studied compounds showed slight stimulatory effects on proliferation of MC3T3-E1 cells, in a dose dependent manner. Additionally, an *in silico* toxicological study of compounds **1** and **2** was performed, and the results indicate that they have a good probability of safe biological intake.

Keywords: Cyclodipeptides, MC3T3-E1 cells, *In silico* toxicological study.

Behind the name "cyclodipeptides", we can unearth a rich world of compounds in terms of structural diversity and biological activity [1]. Cyclodipeptides are secondary metabolites of fungi and plants, or they originate from the marine environment [2]. Structurally, the diversity contained in this class of compounds arises from the type and number of amino acids, many of them completely unusual in nature, size of the macrocycle, and, usually, side chains that adorn the complex macrocycle [1]. Cyclodipeptides contain one or more amino acid(s) replaced by a hydroxy acid, resulting in at least one ester bond in the core ring structure [2]. Furthermore, it has been demonstrated that the ring form of depsipeptides is required for their biological activity [3,4]. The great interest that this class of natural products has elicited in the scientific community is explained by the diverse range of biological activities that they exhibit, including antitumor, antibiotic, antifungal, immunosuppressant, and antiinflammatory. This is supported by intriguing mechanisms of action and attractive molecular architecture [1].

Cyclodipeptides are the simplest among the cyclodipeptides family containing one residue of amino acid and one residue of lactic, α -hydroxyisovaleric or other α -hydroxyacid [5]. They have an ester group and an amide group in the same 6-membered ring [6]. Two cyclodipeptides, 3,6-di-(propan-2-yl)-4-methyl-morpholine-2,5-dione (**1**; C₁₁H₁₉NO₃, M = 213.27) and 3-(2-methylpropyl)-6-(propan-2-yl)-4-methyl-morpholine-2,5-dione (**2**; C₁₂H₂₁NO₃, M = 227.30), were found for the first time in natural products as potential precursors of enniatin B in the pathogenic fungus *Fusarium sporotrichioides* Sherb. This fungus was isolated from the stem of fresh *Hypericum barbatum* Jacq. [6]. The role of compounds **1** and **2** for the producing organism is questionable. For

identification and confirmation, those compounds were synthesized and studied by density functional theory calculations and infrared spectroscopy [6]. Those two 6-(propan-2-yl)-4-methyl-morpholine-2,5-diones were recently evaluated for antimicrobial, immunomodulatory [7], and antioxidant [8] activity, inhibitory activity toward xanthine oxidase (XO) *in vitro* and XO in rat liver homogenate, as well as for antiinflammatory response on human peripheral blood mononuclear cells (PBMCs) [9]. Our data indicate moderate antioxidant [8] and antimicrobial [7] potential of the studied compounds. Furthermore, they were excellent inhibitors of XO and significantly suppressed the activation of nuclear factor κ B [9]. Both, **1** and **2** were confirmed as non-toxic in thymocytes [7] and, therefore, may have promise for use in the treatment of gout and other excessive uric acid production or inflammatory conditions [9].

Hyperuricemia, a potentially harmful condition, favors precipitation of uric acid crystals in joints and tissues, leading to complications such as gout, nephrolithiasis and chronic nephropathy [10]. Gout is an inflammatory arthritis that is the result of the precipitation of serum urate into crystallized deposits of monosodium urate in and around the joints [11]. Also, changes in joints are frequently coupled with changes in surrounding bones such as new bone formation, sclerosis, osteophyte and spur. The finding that bone erosion is strongly associated with all features of new bone formation suggests that loss of bone and formation of new bone may be connected during the joint remodeling process in joints affected by gout. At present, it is unclear whether new bone formation in gout occurs before bone erosion and tophus formation, develops concurrently with bone erosion, or occurs as part of tissue remodeling in response to tophus and erosion [12]. MC3T3-E1 is a

nontransformed continuous osteoblast-like mouse calvarial cell line. Actively growing MC3T3-E1 cells are at the stage of osteoprogenitor cells. They show a typical fibroblastic morphology, possess the capacity to differentiate into osteoblasts [13,14] and frequently are used as model system for studies of cytotoxic and antiproliferative potential of different compounds on bone cells.

Compounds **1** and **2** possess inhibitory activity toward XO [9], a characteristic that recruits them as potential drugs in therapy of gout, clinical manifestation of which often includes changes on bones. In order to obtain more information about the toxicological potential of compounds **1** and **2** on bone cells, the current study was designed to evaluate the effect of these compounds on viability and proliferation of MC3T3-E1 cells. Additionally, an *in silico* toxicological study of two studied 6-(propan-2-yl)-4-methyl-morpholine-2,5-diones was performed.




Figure 1: Chemical structures of the compounds under the study.

Effects of two 6-(propan-2-yl)-4-methyl-morpholine-2,5-diones on viability and proliferation of MC3T3-E1 cells: MC3T3-E1 cells, osteoprogenitor cells derived from mouse calvaria, are a useful tool for examination of cytotoxic and antiproliferative activity of various compounds [14]. Registered change in MC3T3-E1 cell viability ranged from 103.6% to 96.8% after treatment with compound **1**, and from 105.1% to 118.7% (in relation to solvent control) after treatment with compound **2** (Figure 2). The change in MC3T3-E1 cell proliferation ranged from 117.7% to 126.2% and from 102.8% to 119.2% for compounds **1** and **2**, respectively (Figure 3). Treatment with compounds **1** and **2** for 24 h and 72 h in all effective concentrations had no visible effect on cell morphology.

MC3T3-E1 cells grow with a population doubling time of ~ 18 h [14] and it is expected that, under normal conditions, in a period from 24 h until 72 h the cells are divided at least once or more times, which multiply the number of cells. Compounds that reduce cell viability and/or suppress the cell division mechanism will lead to a reduction in the number of cells in culture that will manifest itself as a lower MTT reduction. MTT reduction is associated not only with mitochondria, but also with the cytoplasm and with non-mitochondrial membranes, including the endosome/lysosome compartment and the plasma membrane [17]. Cell death depends, in part, upon mitochondrial dysfunction, which is often characterized by increased production of ROS, increased membrane permeability and eventual release of cell death mediators from mitochondria [18]. We reported earlier [7] that cyclodipeptides evaluated in this research do not decrease a mitochondrial membrane potential (MMP) in rat thymocytes after 24 h of incubation. These results correspond with data obtained from intracellular ROS analysis and cytotoxicity, and may explain, in part, the non-significant ROS production in thymocytes cultured with compounds **1** and **2**. Increasing concentrations of **1** and **2** were able to stimulate proliferation of rat thymocytes [7].

Extracts of dental materials were rated as severely, moderately, slightly or not cytotoxic or antiproliferative where activity relative to control was less than 30%, between 30% and 60%, between 60% and 90% or greater than 90%, respectively [19]. Reverse of the




Figure 2: The effects of different concentrations of compounds **1** and **2** on viability of MC3T3-E1 cells. Results are presented as: mean percentage \pm SD.

* $P < 0.05$; ** $P < 0.01$ compared with control




Figure 3: The effects of different concentrations of compounds **1** and **2** on proliferation of MC3T3-E1 cells. Results are presented as: mean percentage \pm SD.

* $P < 0.05$; ** $P < 0.01$ compared with control

above mentioned classification, positive effect on cell viability or proliferation of tested compounds may be classified as not stimulatory, slightly, moderately and pronounced stimulatory where activity relative to control was between 100% and 110%, between 110% and 140%, between 140% and 170%, and more than 170%, respectively. Compound **1** showed neither cytotoxic nor stimulatory effect on cell viability. Compound **2** showed a slight stimulatory effect on cell viability in a dose dependent manner. Also, in a dose dependent manner compounds **1** and **2** showed a slight stimulatory effect on proliferation of MC3T3-E1 cells.

In silico toxicological study of two 6-(propan-2-yl)-4-methyl-morpholine-2,5-diones: In order to obtain a complete picture of the toxicity of the two studied 6-(propan-2-yl)-4-methyl-morpholine-2,5-diones, we calculated absorption, metabolism and toxicity properties of compounds **1** and **2** using the OSIRIS Property Explorer [20], admetSAR [21] (Table 1) and ACD/I-Lab [22] (Tables 2 and 3).

The data obtained by OSIRIS Property Explorer indicate that the structures of compounds **1** and **2** are supposed to be non-mutagenic, non-tumorigenic, non-irritating and with no reproductive effects. The toxicological properties predicted by admetSAR (Table 1) suggest that two studied compounds might be able to pass through blood-brain barrier and penetrate into the CNS and, therefore, are considered as CNS active compounds. Compounds **1** and **2** were predicted to be capable of being absorbed by the intestine with high probability scores of 0.95 and 0.96, respectively. They are not likely to act as inhibitors of renal organic cation transporter and CYP450 enzymes, but are likely to act as substrates for CYP450 3A4. Additionally, the two studied compounds are supposed to be

weak/non-hERG inhibitors, non-AMES toxic, non-carcinogens, with low fish, *Tetrahymena pyriformis* and honey bee toxicity. While compound **1** is supposed to be ready biodegradable, compound **2** is not. Compounds **1** and **2** are predicted as Category III for risk for acute toxicity, including compounds with LD₅₀ values greater than 500 mg/kg, but less than 5000 mg/kg; and "non-required" rat carcinogenicity class, including non-carcinogenic chemicals (Table 1).

Table 1: *In silico* predicted absorption, metabolism and toxicity properties of compounds **1** and **2** using admetSAR tool.

	Compound 1		Compound 2	
	Result	Probability	Result	Probability
Absorption				
BBB	+	0.8197	+	0.8311
HIA	+	0.9497	+	0.9604
ROC transporter	NI	0.9379	NI	0.8956
Metabolism				
CYP450 2C9 substrate	NS	0.8430	NS	0.8258
CYP450 2D6 substrate	NS	0.8522	NS	0.8158
CYP450 3A4 substrate	S	0.5844	S	0.6170
CYP450 1A2 inhibitor	NI	0.8453	NI	0.8578
CYP450 2C9 inhibitor	NI	0.9354	NI	0.9241
CYP450 2D6 inhibitor	NI	0.9577	NI	0.9461
CYP450 2C19 inhibitor	NI	0.9117	NI	0.8976
CYP450 3A4 inhibitor	NI	0.9845	NI	0.9828
CYP inhibitory promiscuity	Low	0.9699	Low	0.9726
Toxicity				
hERG inhibitor	Weak	0.9958	Weak	0.9939
	NI	0.9719	NI	0.9829
AMES toxicity	–	0.7156	–	0.7573
Carcinogens	–	0.9128	–	0.9053
Fish toxicity	Low	0.9656	Low	0.8434
<i>Tetrahymena pyriformis</i> toxicity	Low	0.9865	Low	0.9600
Honey bee toxicity	Low	0.6351	Low	0.6684
Biodegradation	Ready biodegradable	0.5259	Not ready biodegradable	0.6323
Acute oral toxicity	III	0.7319	III	0.7013
Carcinogenicity (Three-class)	Non-required	0.6836	Non-required	0.6797

*BBB – blood-brain barrier; HIA – human intestinal absorption; ROC – Renal organic cation; NS – non-substrate; S – substrate; NI – non-inhibitor; hERG – human ether-a-go-go-related gene.

Table 2: Health effects of compounds **1** and **2** calculated using ACD/I-Lab.

Health effects	Compound 1	Compound 2
Gastrointestinal system	0.48	0.59
Lungs	0.28	0.26
Cardiovascular system	0.53	0.52
Liver	0.1	0.09
Blood	0.11	0.12
Kidney	0.15	0.15

*Values around 0.00 indicate low, around 0.50 moderate, and around 1.00 high probabilities of health effects on certain organ systems.

According to the results obtained using ACD/I-Lab, compounds **1** and **2** have low probability of health effects on liver, blood, kidney and lungs, and moderate probability of health effects on cardiovascular and gastrointestinal system (Table 2). Using ACD/I-Lab software we predicted LD₅₀ values in mg/kg after intraperitoneal, oral, intravenous and subcutaneous administration to mice, and after intraperitoneal and oral administration to rats (Table 3). The data indicate that compounds **1** and **2** are slightly toxic having oral LD₅₀ values between 500 and 5000 mg/kg, and therefore are predicted as Category III for acute toxicity.

In conclusion, both tested compounds show a slightly dose dependent stimulatory effect on MC3T3-E1 cell proliferation. Summarizing the physico-chemical properties of cyclodipeptides, we concluded in our recent review [5] that they obey the Lipinski "Rule of five" and meet all criteria for good solubility and permeability. The results of *in silico* studies shown in

Table 3: LD₅₀ values of compounds **1** and **2** calculated using ACD/I-Lab.

Species/ Administration route	LD ₅₀ (mg/kg)		Reliability (RI)
	Compound 1	Compound 2	
Mouse/Intraperitoneal	550	550	Borderline (0.47)
Mouse/Oral	2000	2100	Moderate (0.58)
Mouse/Intravenous	78	73	Not reliable (0.29)
Mouse/Subcutaneous	510	490	Borderline (0.42)
Rat/Intraperitoneal	860	850	Borderline (0.5)
Rat/Oral	2700	2800	Borderline (0.3)

Tables 1-3 indicate a good probability of compounds **1** and **2** being biologically safe for intake and make them promising leads for future development of safe and efficient agents. So, two studied 6-(propan-2-yl)-4-methyl-morpholine-2,5-diones have promise to be used in the treatment of gout and other excessive uric acid production or inflammatory conditions.

Experimental

Cell culture: MC3T3-E1 cells were cultivated in Dulbecco's modified Eagle's medium (DMEM) (PAA Laboratories GmbH), supplemented with 10% fetal calf serum (FCS) (Gibco, Great Britain), 100 IU/mL penicillin, 100 µg/mL streptomycin and 4 mM glutamine. Cells were incubated in a humidified chamber at 37°C containing 5% CO₂.

Test compounds: The synthesis of the 2 cyclodipeptides used in the study, **1** and **2**, was performed via *N*-(α -bromoacyl)- α -amino acids, as described in our previous study [6].

The effects of two 6-(propan-2-yl)-4-methyl-morpholine-2,5-diones on the proliferation and viability of MC3T3-E1 cells were tested *in vitro* using MTT assay according to a slightly modified protocol [15].

Cell viability assay: Exponentially growing cells MC3T3-E1, (2×10^4 cells in 0.1 mL of medium/well) were seeded on 96-well plates in triplicate. Cells were incubated overnight in an atmosphere saturated with aqueous vapor, at 37°C with 5% CO₂. After 24 h, medium with various concentrations of ethanolic solutions of compounds **1** and **2** in a final volume of 0.1 mL was added to the wells. Effective concentrations of test compounds **1** and **2** were 0.1, 1, and 10 µg/mL.

At the same time, medium with an appropriate amount of 96% ethanol was added to control samples. Cells were incubated for 24 h. After the incubation the supernatant was removed and 0.1 mL of phosphate buffered saline (PBS) and 0.02 mL of 3-(4,5-dimethylthiazol-2-yl)-2,5-diphenyl-tetrazoliumbromide (MTT) (Carl Roth, Germany), previously diluted in PBS, were added to each well and incubated for 2.5 h. After the estimated time, the reduced formazan product was dissolved in 0.1 mL of isopropanol per well. The absorbance was read at 540 nm on a multichannel spectrophotometer (Multiskan Ascent N°354, Thermo Labsystems, Finland).

Cell proliferation assay: Exponentially growing cells MC3T3-E1 (0.5×10^4 cells in 0.1 mL of medium/well) were seeded into 96-well plates in triplicate. Cells were incubated overnight in an atmosphere saturated with aqueous vapor, at 37°C with 5% CO₂. After 24 h, medium with various concentrations of ethanolic solutions of compounds **1** and **2** in a final volume of 0.1 mL was added to the wells. Effective concentrations of test compounds **1** and **2** were 0.1, 1, and 10 µg/mL. At the same time, medium with an appropriate amount of 96% ethanol was added to control samples. Cells were incubated for 72 h. After the estimated time, MTT assay was carried out in the same way as in the cell viability assay.

Data analysis: Absorbance of the control was regarded as 100% and values (%) of the different concentrations of test compounds were calculated as follows:

absorbance of the compound/absorbance of the control $\times 100\%$ = percent of the compound in comparison with a control.

Results are presented as: mean \pm SD. Statistical differences between different concentrations of the evaluated compounds in comparison with control were analyzed in SPSS 17.0 with Student's t-test and considered significant at $*P < 0.05$ and $**P < 0.01$.

Cell morphology: Cells that were used are adherent, and the possible toxic effect of the test compounds was evaluated through

the possible change of their morphological characteristics. The normal morphology of MC3T3-E1 pre-osteoblast cells is polygonal [16]. Alteration of its phenotype was considered as a sign of a toxic effect. Cell morphological characteristics were observed after 24 h and 72 h of incubation using an invert microscope (AxioObserver.Z1 Carl Zeiss, Germany).

Acknowledgments - The financial support of this work by the Ministry of Education and Science of the Republic of Serbia (Grant No.172044) and National Science Fund of Bulgaria (young researchers project DMU-03/66) is gratefully acknowledged.

References

- [1] Sarabia F, Chammaa S, Ruiz AS, Ortiz LM, Herrera FL. (2004) Chemistry and biology of cyclic depsipeptides of medicinal and biological interest. *Current Medicinal Chemistry*, **11**, 1309-1332.
- [2] Lemmens-Gruber R, Kamyar MR, Dornetshuber R. (2009) Cyclodepsipeptides - potential drugs and lead compounds in the drug development process. *Current Medicinal Chemistry*, **16**, 1122-1137.
- [3] Hamann MT, Otto CS, Scheue PJ, Dunbar DC. (1996) Kahalalides: bioactive peptides from a marine mollusk *Elysia rufescens* and its algal diet *Bryopsis* sp. *The Journal of Organic Chemistry*, **61**, 6594-6600.
- [4] Isaka M, Palasarn S, Lapanun S, Sriklung K. (2007) Paecilodepsipeptide A, an antimalarial and antitumor cyclohexadepsipeptide from the insect pathogenic fungus *Paecilomyces cinnamomeus* BCC 9616. *Journal of Natural Products*, **70**, 675-678.
- [5] Smelcerovic A, Dzodic P, Pavlovic V, Cherneva E, Yancheva D. (2014) Cyclodidepsipeptides with a promising scaffold in medicinal chemistry. *Amino Acids*, **46**, 825-840.
- [6] Smelcerovic A, Yancheva D, Cherneva E, Petronijevic Z, Lamshoeft M, Herebian D. (2011) Identification and synthesis of three cyclodidepsipeptides as potential precursors of enniatin B in *Fusarium sporotrichioides*. *Journal of Molecular Structure*, **985**, 397-402.
- [7] Pavlovic V, Djordjevic A, Cherneva E, Yancheva D, Smelcerovic A. (2012) Stimulatory effect on rat thymocytes proliferation and antimicrobial activity of two 6-(propan-2-yl)-4-methyl-morpholine-2,5-diones. *Food and Chemical Toxicology*, **50**, 761-766.
- [8] Stankov-Jovanovic V, Tabet JC, Dzodic P, Daskalova L, Cherneva E, Yancheva D, Smelcerovic A. (2012) *In vitro* antioxidant activity of two 6-(propan-2-yl)-4-methyl-morpholine-2,5-diones. *Acta Chimica Slovenica*, **59**, 939-943.
- [9] Smelcerovic A, Rangelov M, Smelcerovic Z, Veljkovic A, Cherneva E, Yancheva D, Nikolic GM, Petronijevic Z, Kocic G. (2013) Two 6-(propan-2-yl)-4-methyl-morpholine-2,5-diones as new non-purine xanthine oxidase inhibitors and anti-inflammatory agents. *Food and Chemical Toxicology*, **55**, 493-497.
- [10] Grassi D, Ferri L, Desideri G. (2013) Chronic hyperuricemia, uric acid deposit and cardiovascular risk. *Current Pharmaceutical Design*, **19**, 2432-2438.
- [11] Dalbeth N, Lauterio T, Wolfe H. (2014) Mechanism of action of colchicine in the treatment of gout. *Clinical Therapeutics*, **36**, 1465-1479.
- [12] Dalbeth N, Milligan A, Doyle AJ, Clark B, McQueen FM. (2012) Characterisation of new bone formation in gout: a quantitative site-by-site analysis using plain radiography and computed tomography. *Arthritis Research and Therapy*, **14**, R165.
- [13] Kodama H, Amagai Y, Sudo H, Kasai S, Yamamoto S. (1981) Establishment of a clonal osteogenic cell line from new born mouse calvaria. *Japanese Journal of Oral Biology*, **23**, 899-901.
- [14] Sudo H, Kodama HA, Amagai Y, Yamamoto S, Kasai S. (1983) *In vitro* differentiation and calcification in a new clonal osteogenic cell line derived from newborn mouse calvaria. *The Journal of Cell Biology*, **96**, 191-198.
- [15] Mosmann T. (1983) Rapid colorimetric assay for cellular growth and survival: application to proliferation and cytotoxicity assays. *Journal of Immunological Methods*, **65**, 55-63.
- [16] Horikawa A, Okada K, Sato K, Sato M. (2000) Morphological changes in osteoblastic cells (MC3T3-E1) due to fluid shear stress: cellular damage by prolonged application of fluid shear stress. *The Tohoku Journal of Experimental Medicine*, **191**, 127-137.
- [17] Berridge MV, Herst PM, Tan AS. (2005) Tetrazolium dyes as tools in cell biology: new insights into their cellular reduction. *Biotechnology Annual Review*, **11**, 127-152.
- [18] Exline MC, Crouser ED. (2008) Mitochondrial mechanisms of sepsis induced organ failure. *Frontiers in Bioscience*, **13**, 5030-5041.
- [19] Lönnroth EC, Dahl JE. (2003) Cytotoxicity of liquids and powders of chemically different dental materials evaluated using dimethylthiazol diphenyltetrazolium and neutral red tests. *Acta Odontologica*, **61**, 52-56.
- [20] OSIRIS Property Explorer, <http://www.organic-chemistry.org/prog/peo/>
- [21] admetSAR, <http://lmmd.ecust.edu.cn:8000/predict/>
- [22] ACD-I/Lab, <https://ilab.acdlabs.com/>

Red Maple (<i>Acer rubrum</i>) Aerial Parts as a Source of Bioactive Phenolics Yan Zhang, Hang Ma, Tao Yuan and Navindra P. Seeram	1409
Changes in the Content of the Glycosides, Aglycons and their Possible Precursors of <i>Rhodiola rosea</i> during the Vegetation Period Iman Mirmazloun, Márta Ladányi and Zsuzsanna György	1413
Biocompounds Attenuating the Development of Obesity and Insulin Resistance Produced by a High-fat Sucrose Diet Usune Etxeberria, Ana Laura de la Garza, J. Alfredo Martínez and Fermín I. Milagro	1417
A Novel C21 Cyclopentenone Derivative from <i>Cipadessa cinerascens</i> Jing Zhang, Yao-Wen Chang, Chun-Mao Yuan, Fan Zhang, Ying-Tong Di and Xiao-Jiang Hao	1421
Effects on MC3T3-E1 Cells and <i>In silico</i> Toxicological Study of Two 6-(Propan-2-yl)-4-methyl-morpholine-2,5-diones Marija Vukelić-Nikolić, Ana Kolarević, Katarina Tomović, Denitsa Yancheva, Emiliya Cherneva, Stevo Najman, and Andrija Šmelcerović	1423
Release of Antioxidant Peptides from the Body Wall Proteins of the Sea Cucumber <i>Isostichopus fuscus</i> Arisaí C. Hernández-Sámano and Blanca Hernández-Ledesma	1427
Bioactivity-guided Separation of the Active Compounds in <i>Acacia pennata</i> Responsible for the Prevention of Alzheimer's Disease Pattamapan Lomarat, Sirirat Chancharunee, Natthinee Anantachoke, Worawan Kitphati, Kittisak Sripha and Nuntavan Bunyapraphatsara	1431
Development and Validation of LC-MS/MS Method for Quantitative Determination of Adenosine, Guanosine, Xanthine and Uric acid in Widely Consumed Vegetables in Thailand Narisa Rukdee, Piyanuch Rojsanga and Chutima Matayatsuk Pechkrajang	1435
Chemical Composition of the Essential Oil from <i>Chaerophyllum temulum</i> (Apiaceae) Jelena G. Stamenković, Gordana S. Stojanović, Ivana R. Radojković, Goran M. Petrović and Bojan K. Zlatković	1439
Comparison of Essential Oils Obtained from Different Extraction Techniques as an Aid in Identifying Aroma Significant Compounds of Nutmeg (<i>Myristica fragrans</i>) Suchandra Chatterjee, Sumit Gupta and Prasad. S. Variyar	1443
Chemical Variability of the Essential Oil Isolated from Aerial Parts of <i>Tetraclinis articulata</i> from North-Western Algeria Maghnia Boussaïd, Chahrazed Bekhechi, Fawzia Beddou, Daoudi Chabane Sari, Ange Bighelli, Joseph Casanova and Félix Tomi	1447
Composition and Bioactivities of an (<i>E</i>)-β-Farnesene Chemotype of Chamomile (<i>Matricaria chamomilla</i>) Essential Oil from Nepal Prabodh Satyal, Samon Shrestha and William N. Setzer	1453
Chemical Composition of the Essential Oil from <i>Croton oblongifolius</i> and its Antibacterial Activity against <i>Propionibacterium acnes</i> Sirivan Athikomkulchai, Sarin Tadtong, Nijsiri Ruangrungrasi and Tapanee Hongratanaworakit	1459
Composition, <i>in vitro</i> Anti-inflammatory, Antioxidant and Antimicrobial Activities of Essential Oils from Leaf and Twig Parts of <i>Cupressus cashmeriana</i> Yu-Chang Su, Kuan-Ping Hsu, Kuo-Feng Hua and Chen-Lung Ho	1461
Antioxidant and Anticholinesterase Activities of Essential Oils of <i>Cinnamomum griffithii</i> and <i>C. macrocarpum</i> Wan Mohd Nuzul Hakimi Wan Salleh, Farediah Ahmad and Khong Heng Yen	1465
Essential Oil Composition and Antigermination Activity of <i>Artemisia dracunculus</i> (Tarragon) Daniele Fraternali, Guido Flamini and Donata Ricci	1469
<i>In Vitro</i> Activity of Twenty Commercially Available, Plant-Derived Essential Oils against Selected Dermatophyte Species Simona Nardoni, Silvia Giovanelli, Luisa Pistelli, Linda Mugnaini, Greta Profili, Francesca Pisseri and Francesca Mancianti	1473
Intracerebral Distribution of α-Pinene and the Anxiolytic-like Effect in Mice Following Inhaled Administration of Essential Oil from <i>Chamaecyparis obtusa</i> Hikaru Kasuya, Sayuka Iida, Kurumi Ono, Tadaaki Satou and Kazuo Koike	1479
<u>Accounts/Reviews</u>	
A Comprehensive Review of the Cosmeceutical Benefits of <i>Vanda</i> Species (Orchidaceae) Hazrina Hadi, Syarifah Nazira Said Razali and Ammar Ihsan Awadh	1483
Genotoxicity and Antigenotoxicity Studies of Traditional Medicinal Plants: How Informative and Accurate are the Results? Luc Verschaeve	1489

Natural Product Communications

2015

Volume 10, Number 8

Contents

Original Paper

- Isolation of Monovalerianester A, an Inhibitor of Fat Accumulation, from *Valeriana fauriei***
Keiji Yuki, Mariko Ikeda, Shosuke Yoshida, Osamu Ohno, Kiyotake Suenaga, Kaoru Yamada, Daisuke Uemura and Kenji Miyamoto 1333
- Arthropod Deterrents from *Artemisia pallens* (Davana Oil) Components**
Ganga V. Bhagavathy, Glory M Velazquez Nieves, Meiling Z. Webb and Kamlesh R. Chauhan 1335
- Stereoselective Synthesis of 2,15-Dihydroxycalamenene and 2-Methoxycalamenene. Determination of the Configuration of Natural 2,15-Dihydroxycalamenene**
Stefano Serra 1337
- Sesquiterpenes from the Vietnamese Marine Sponge *Dysidea fragilis***
Nguyen Thi Cuc, Hoang Le Tuan Anh, Dan Thi Thuy Hang, Nguyen Xuan Nhiem, Nguyen Hai Dang, Nguyen Hoai Nam, Pham Hai Yen, Do Cong Thung, Vu Kim Thu, Chau Van Minh and Phan Van Kiem 1341
- Vibrational Circular Dichroism Absolute Configuration of 9,12-Cyclomulin-13-ol, a Diterpene from *Azorella* and *Laretia* Species**
Marcelo A. Muñoz, Aurelio San-Martin and Pedro Joseph-Nathan 1343
- Ultrafine Betulin Formulation with Biocompatible Carriers Exhibiting Improved Dissolution Rate**
Svetlana A. Myz, Tatyana P. Shakhtshneider, Mikhail A. Mikhailenko, Andrey G. Ogienko, Ekaterina G. Bogdanova, Anna A. Ogienko, Svetlana A. Kuznetsova, Elena V. Boldyreva and Vladimir V. Boldyrev 1345
- Bio-assay Guided Isolation of Anti-cancer Compounds from *Anthocephalus cadamba* Bark**
Deepak Kumar, Chilukuri Tejaswi, Saiprasanna Rasamalla, Sumana Mallick and Bikas C Pal 1349
- Damarane-type Saponins from *Gynostemma longipes* and their Cytotoxic Activity**
Pham Tuan Anh, Pham Thanh Ky, Nguyen Thi Cuc, Nguyen Xuan Nhiem, Pham Hai Yen, Tran Minh Ngoc, Hoang Le Tuan Anh, Bui Huu Tai, Do Thi Trang, Chau Van Minh and Phan Van Kiem 1351
- A New C₂₃ Steroid from the Venom of *Bufo bufo gargarizans***
Shi-Lin Luo, Hai-Yan Tian, Jun-Shan Liu, Ying Wang and Wen-Cai Ye 1353
- (-)-Pentylsedinine, a New Alkaloid from the Leaves of *Lobelia tupa* with Agonist Activity at Nicotinic Acetylcholine Receptor**
Cristian Paz, José Becerra, Mario Silva, Viviana Burgos, Matthias Heydenreich, Bernd Schmidt, Thu Tran and Irina Vetter 1355
- Furoquinoline Alkaloids from the Leaves of *Evodia lepta* as Potential Cholinesterase Inhibitors and their Molecular Docking**
Jirapast Sichaem, Thanawan Rojpitikul, Pattara Sawasdee, Kiattisak Lugsanangarm and Santi Tip-pyang 1359
- Asperginine, an Unprecedented Alkaloid from the Marine-derived Fungus *Aspergillus* sp.**
Pinmei Wang, Shizhe Zhao, Ying Liu, Wanjing Ding, Feng Qiu and Jinzhong Xu 1363
- Effect of Quercetin on Cell Cycle and Cyclin Expression in Ovarian Carcinoma and Osteosarcoma Cell Lines**
Daniela Catanzaro, Eugenio Ragazzi, Caterina Vianello, Laura Caparrotta and Monica Montopoli 1365
- Anti-*Helicobacter pylori* Activity of Four *Alchemilla* Species (Rosaceae)**
Marija Krivokuća, Marjan Niketić, Marina Milenković, Nataša Golić, Carla Masia, Maria Maddalena Scaltrito, Francesca Sisto and Tatjana Kundaković 1369
- Peracylated Glucosyl Kaempferols from *Pasania dodoniifolia* Leaf**
Chi-Chih Chang and Shoei-Sheng Lee 1373
- 6-Methoxyflavonol Glycosides with *In Vitro* Hepatoprotective Activity from *Chenopodium bonus-henricus* Roots**
Zlatina Kokanova-Nedialkova, Magdalena Kondeva-Burdina, Dimitrina Zheleva-Dimitrova, Virginia Tzankova, Stefan Nikolov, Jörg Heilmann and Paraskev T. Nedialkov 1377
- Flavonol Glycosides from the Leaves of *Allium macrostemon***
Risa Nakane and Tsukasa Iwashina 1381
- HPLC Plasma Assay of a Novel Anti-MRSA Compound, Kaempferol-3-O-Alpha-L-(2'',3''-di-p-coumaroyl)rhamnoside, from Sycamore Leaves**
Yiguan Zhang, Frederick Valeriote, Kenneth Swartz, Ben Chen, Mark T. Hamann, Douglas L. Rodenburg, James D. McChesney and Jiajiu Shaw 1383
- Biological Activity of *Dolichandrone serrulata* Flowers and Their Active Components**
Phanida Phanthong, Noppawan Phumala Morales, Sirirat Chancharunee, Supachoke Mangmool, Natthinee Anantachoke and Nuntavan Bunyapraphatsara 1387
- Prenylhydroquinone-Derived Secondary Metabolites from Cultures of the Basidiomycete *Lentinus similis* BCC 52578**
Masahiko Isaka, Somporn Palasarn, Malipan Sappan, Kitlada Srichomthong, Samantha C. Karunarathna and Kevin D. Hyde 1391
- Effects of Thymoquinone on the Pharmacokinetics and Pharmacodynamics of Glibenclamide in a Rat Model**
Ajaz Ahmad, Rao Muzaffar A. Khan, Khalid M. Alkharfy, Mohammad Raish, Fahad I. Al-Jenoobi, and Abdullah M. Al-Mohizea 1395
- Antioxidant and Antiinflammatory Compounds in Nutmeg (*Myristica fragrans*) Pericarp as Determined by *in vitro* Assays**
Chuan-Rui Zhang, Ettannil Jayashree, Paramasivam Suresh Kumar and Muraleedharan G. Nair 1399
- In Vitro* Safety/Protection Assessment of Resveratrol and Pterostilbene in a Human Hepatoma Cell Line (HepG2)**
Germana Lombardi, Samuele Vannini, Francesca Blasi, Maria Carla Marcotullio, Luca Dominici, Milena Villarini, Lina Cossignani and Massimo Moretti 1403

Continued inside backcover